STRUCTURE AC	CTIVITY TEA	M REPORT	ver. 0	4/98		
Case #:	L-08-0037		DCN:			
SAT Date:	11/2/2007		SAT Ch	air:	<del>-11/02/2007</del>	- V. NAbholz
Submitter:						
Chemical Name:						
						,
CAS RN:			Trade Na	me:		
Structure	<u> </u>				. v	
						23
						2007
						.4
						C)
						en e e e e e e e e e e e e e e e e e e
						<u>.</u>
Molecular Formula:						100
Molecular Wt.		WT%<500:			WT%<1000:	
MP:		BP:			Eq. Wt:	/
H2O Sol (g/L):		Dispe	ersible	V.P.		<0.0325
Max. Prod. Volume (kg/	/yr):			hysical State:		Solid
USE:						
Emulsifier for polymeriza	tion reactions					
There is 1 reference in fil	e CA on STN.	Case Role		Related Ca	26	
Related Cas	e Numbers	Case Roin	3	Relateu C	<b>a</b> 5	
			Ţ			
Focus Date: NO	v 19 2007	Results:	Den	1 of 9		4
			Page	of <b>9</b>		

STRUCTURE ACTIVITY TEAM REPORT 02 November 2007 CASE NUMBER: L08-0037 RELATED CASES: CONCLUSIONS/DISCUSSIONS TYPE OF CONCERN: **HEALTH** ECOTOX 3 LEVEL: 3 KEYWORDS: IRR-S, LIVER, REPRO, DEVEL, IMMUNO, ONCO, AQUATOX-A,C, LUNGS SUMMARY OF ASSESSMENT: FATE: solid with mp = sublimes at 204 C. (M) log Kow = 2.0 (EPI)S = dispersible (P)vp < 0.032 mm Hg or torr at 25 °C (P)</pre> bp = sublimes at 204 C (M) H < 1.0E-8 (P) log Koc = 3.1 (P)log fish BCF = 0.50 (P)sorption to sludge = moderate submitted test data for aerobic biodegradation were: 22% biodegradation in 28 d via CO2 (Closed bottle [OECD301D]), thus, not readily biodegradable POTW removal = 0% time for complete ultimate aerobic biodegradation > months sorption to soils and sediments = moderate PBT Potential: P2B1T3 \*CEB FATE: migration to ground water = moderate HEALTH: Absorption poor thru skin, good thru lungs, and moderate thru GI tract based on analogs; submitted test data with this PMN were: rat acute oral LD100 = 2.0 g/kg with systemic toxic signs, with NOEL = 300 mg/kg;moderate skin irritation in rabbits; severe eye irritation in rabbits; Ames test was negative; E. coli test was negative; no skin sensitization in mice up to 50% ai (LLNA);

By analogy to PFOA, there is concern for reproductive and developmental effects, liver effects, immunotoxicity, and a marginal concern for oncogenicity. In the assessment conducted by the state of West Virginia with input from EPA, the RfD for oral exposure was determined to be 0.004 mg/kg/day and the RfC for inhalation exposure was determined to be 1 ug/m3. Some of the key findings for non-cancer assessment are for PFOA and/or its salts:

and the second control of the second second

its salts:
90-day dietary study in male rats - NOAEL = 0.47 mg/kg/day based on liver effects at 1.44 mg/kg/day. The benchmark dose was 1.3 mg/kg/day;

two-generation reproduction study in rats, gavage - LOAEL = 1
mg/kg/day based on increased liver and kidney weights in P and F1
generations;

2-year dietary study in rats - NOAEL = 1.3 mg/kg/day in males based on increased liver weight, hepatic cystoid degeneration, increased ALT enzyme activity, testicular vascular mineralization and LOAEL = 1.6 mg/kg/day in females based on an increase in the incidence of ovarian stromal tubular hyperplasia; the benchmark dose was 0.73 mg/kg/day based on liver effects in males; 26-week oral study in cynomolgus monkeys - LOAEL is from 3 to 10 mg/kg/day based on 30% increased absolute liver weight with no NOAEL;


2-week dermal study in rats - increased liver weight and liver pathology;

2-week inhalation study in male rats - liver effects at 7.6 mg/m3; severe toxicity including the death of one animal at 84 mg/m3, increased lung and testes weight at 84 mg/m3; inhalation developmental toxicity study in rats - maternal deaths, reduced maternal weights, reduced fetal weights at 25 mg/m3;

oral developmental toxicity in rabbits - increase in skeletal variations at 50 mg/kg.

high concern for toxicity
\*CEB HEALTH: Exposures to humans! enhalation, dermal, ingestion, and drinking water;

- moderno (o mumimos: immagation contri



Predictions are based on SAR-test data for PFOA; SAR chemical class = PFOA; pH7; effective concentrations based on 100% active ingredients and mean measured concentrations; DW hardness < 150.0 mg/L as CaCO3; and DW TOC <2.0 mg/L;

moderate concern for aquatic toxicity;

high concern for reproductive toxicity to birds and wild mammals due to chronic toxicity observed in mammals;

assessment factor

concern concentration = 0.050 mg/L (ppm)

\*CEB ECOTOX: All releases to the environment with CC = 50 ppb

SAT Co-chair:

aria i aranga da in manana Ta.

## **GTOX Report**


PMN No. L-08-0037 S/A Name of Analog S	lo.	Rcvd: 10/24/07	OECD Incomplet	ID: Rec# 8 : 2  Reviewer
		with activation	without activation	Positive Strains
Salmonella Assay:		N	N	
Chromosomal Aberration	CHO: CHL: V79:			
E. coli Reverse Mutation:	-	N	N	
Mouse Micronucleus Assay:		Route:		
Rat Hepatocytes Unschedule	d DNA S	ynthesis:		
Other GTOX Results A Salmonella assay was also	o provide	for the analog		
ЕСОТОХ:		MM2		
Fate:		Ready B	iodegradability, (pg. 72-	89)
				ppe.
WS/Log P:	W	S => 87% w/w @ 20	)C (M, pg. 43), LogP = -0	.403 (M, pg. 43).

## **Toxicology Report**

PMN No. L- <b>08-0</b> (	CAS No.	Rcvd: 10/24/2007	OECD Incomplete		ID: Rec# 8	: 5
S/A S	Name of Analog				Reviewer	Study#: <b>5</b>
Study Ty EIRR	pe	Species RABB	Sex Route NS EYES			
	stance Description olid, purity: 96.4-10	0%, impurity: H substitut	ed 3.6mol%			
Test Con	ditions					
each of superfu	three enucleated e ision chamber. Tw	Test Conditions (Dose re yes which had been main o control eyes were treat pre-enucleation, post eq	ntained at a tempera ed with saline solut	iture of 32 C + ion. Measurer	/- 1.5 C within nents of corne	a al
		ute observations 23.4%, erial was considered to h				
				<i>ې</i> ېژ <i>ىر</i> ر		

## **Toxicology Report**

PMN No. CAS No.  L-08-0037  S/A Name of Analog  S	Revd: 10/24/2007	OECD Incomplete		ID: Rec# 8 Reviewer EC	: 7 Study#: 7
Study Type DSEN	Species MICE	Sex Route F DERM	!		
Test Substance Description white solid, purity: 96.4-100%,	impurity: H substitut	ed 3.6mol%, carrier	: dimethyl forma	míde	- van
Test Conditions Study duration: 6 days; Strain: Controls: positive control (a-Hregimen): The study was cond Following a preliminary screen surface of the ear for 3 consec 3H-methyl thymidine and then indices were determined.	exylcinnamaldehyde, ucted according to O ling, mice were expo- utive days. Five days	, Tech, 85%), 1/4; Do PECD 429 (skin sens sed to 25 uL/ear of t s following first app	ose Level: 25 ul; sitization: Local I the test substan olication, mice w	Test Condition  _ymph Node    ce on the dorse  ere injected w	ons (Dose Assay). sal
} 1				,	
Results Stimulation indices were 1.36, respectively. The test material					st. ************************************
			pýk.		
1 1 1 1					} ;



		<del></del>	
	ATTENDEES	SIGNATURE	
CHEM	MISTRY		
	Paul Bickart Diana Darling Rich Engler Greg Fritz Daniel Lin Kathy Schechter	Kalty Scheelite	
ENVI	RONMENTAL FATE		
	Bob Boethling Wen-Hsiung Lee Laurence Libelo David Lynch Andy Mamantov	Only Marices	Aw
HEAL	тн		
	Katherine Anitole Michael Cimino Steve Cragg Leonard Keifer David Lai Jim Murphy Deborah Norris Ronald Ward Yin Tak Woo	Mento Cicio	
ENVI	RONMENTAL EFFECTS		
	Gordon Cash Jince Nabholz Maggie Wilson	МЕМО	
SAT C	CHAIR/OTHER		
V	Rebecca Jones Geonard Keifer Vince Nabholz Vim Kwiat	Mulloz	14 <sup>1</sup> .